



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

Zeros of the Hurwitz zeta function in the interval (0,1)

Schipani, D

Abstract: We first give a condition on the parameters s, w under which the Hurwitz zeta function $\zeta(s, w)$ has no zeros and is actually negative. As a corollary we derive that it is nonzero for $w \geq 1$ and $s \in (0, 1)$ and, as a particular instance, the known result that the classical zeta function has no zeros in $(0, 1)$.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-55175>

Journal Article

Accepted Version

Originally published at:

Schipani, D (2011). Zeros of the Hurwitz zeta function in the interval (0,1). *Journal of Combinatorics and Number Theory*, 3(1):71-74.

Zeros of the Hurwitz zeta function in the interval $(0, 1)^*$

Davide Schipani
Institute of Mathematics
University of Zürich-
IBM Zürich Research Lab
davide.schipani(at)math.uzh.ch

June 3, 2011

Abstract

We first give a condition on the parameters s, w under which the Hurwitz zeta function $\zeta(s, w)$ has no zeros and is actually negative. As a corollary we derive that it is nonzero for $w \geq 1$ and $s \in (0, 1)$ and, as a particular instance, the known result that the classical zeta function has no zeros in $(0, 1)$.

1 Statement and proof of the results

The Hurwitz zeta function is classically defined for $\Re(s) > 1$ as

$$\zeta(s, w) \doteq \sum_{n=0}^{\infty} (n+w)^{-s},$$

with w being a positive real number, and it can be continued analytically to the whole s -plane, except for a pole in 1 (see e.g. [2, Proposition 9.6.6]). Sometimes the definition is extended by letting w be a complex number, while in other situations w is only restricted to be a real number in $(0, 1]$. Notice in fact the following relation:

$$\zeta(s, w) = \zeta(s, w+1) + w^{-s}$$

which follows by considering $n+1$ instead of n .

We set as usual $\sigma = \Re(s)$. The following theorem is meant to add a new result to what is already known about its zeros (see e.g. [4]).

*Research supported by Swiss National Science Foundation Grant no. 107887. Support has also been given over time by Scuola Normale Superiore of Pisa, University of Zürich and IBM Zürich Research Lab.

Theorem 1.1. *Suppose that $1 - w \leq \sigma$. Then $\zeta(s, w)$ is negative and in particular nonzero for $s \in (0, 1)$.*

Proof. As a first step we derive a representation for the Hurwitz zeta function through the Euler-Maclaurin summation formula, in analogy with the one derived for the classical Riemann zeta function (see e.g [3, chapter 6]).

Namely, the following Euler-Maclaurin summation formula

$$\sum_{n=N}^M f(n) = \int_N^M f(x)dx + \frac{1}{2}[f(M) + f(N)] + \frac{B_2}{2}f'(x)|_N^M + \frac{B_4}{4!}f^{(3)}(x)|_N^M \dots,$$

where B_2, B_4, \dots are Bernoulli numbers, is applied to $f(n) = (n + w)^{-s}$ letting M tend to ∞ , to obtain, for $\Re(s) > 1$,

$$\zeta(s, w) = \sum_{n=0}^{N-1} (n + w)^{-s} + \frac{(N + w)^{1-s}}{s - 1} + \frac{1}{2}(N + w)^{-s} + O((N + w)^{-\sigma-1}).$$

One then argues that this formula is in fact true for all $\mathbb{C} \setminus \{1\}$, as explained for the zeta function ([3, section 6.4]), by considering an expression for the rest in integral form and its correspondent halfplane of convergence (see also [2, Proposition 9.6.7]).

The formula above implies that, if we restrict to $\Re(s) > 0$,

$$\zeta(s, w) = \lim_{N \rightarrow \infty} \left(\sum_{n=0}^N (n + w)^{-s} - \frac{(N + w)^{1-s}}{1 - s} \right).$$

Now, by the triangular inequality,

$$\left| \sum_{n=0}^N (n + w)^{-s} \right| \leq \sum_{n=0}^N (n + w)^{-\sigma} < w^{-\sigma} + \int_0^N (x + w)^{-\sigma} dx = w^{-\sigma} + \frac{(N + w)^{1-\sigma}}{1 - \sigma} - \frac{w^{1-\sigma}}{1 - \sigma}$$

(in the case of real s , which is what will interest us now, the first comparison is in fact an equality)

which we write as

$$\frac{(N + w)^{1-\sigma}}{1 - \sigma} + \frac{(1 - \sigma)w^{-\sigma} - w^{1-\sigma}}{1 - \sigma}.$$

Suppose first that $(1 - \sigma)w^{-\sigma} < w^{1-\sigma}$, which is the same as $1 - w < \sigma$. Since $1 - \sigma$ is positive for $\sigma < 1$, then the expression above is less than

$$\frac{(N + w)^{1-\sigma}}{1 - \sigma} - a,$$

where a is a fixed positive quantity.

Therefore, in $(0, 1)$ and for every N , $\left| \sum_{n=0}^N (n + w)^{-s} \right|$ which is also $\sum_{n=0}^N (n + w)^{-s}$ or $\sum_{n=0}^N (n + w)^{-\sigma}$ is less than $\left| \frac{(N + w)^{1-s}}{1-s} \right| = \frac{(N + w)^{1-s}}{1-s} = \frac{(N + w)^{1-\sigma}}{1-\sigma}$ by more than a fixed positive quantity. Then in the limit the difference is nonzero, and in fact negative.

For the case $1 - w = \sigma$, notice that we would not be able to conclude if, as N goes to ∞ , the difference between $\sum_{n=0}^N (n + w)^{-\sigma}$ and $w^{-\sigma} + \int_0^N (x + w)^{-\sigma} dx$ could decrease and tend to 0. But this is not the case, since it is actually increasing, as the approximation with the integral is accumulating error as N gets bigger. The thesis then follows also in this scenario. \square

Corollary 1.2. *Suppose $w \geq 1$. Then $\zeta(s, w)$ has no zeros for $s \in (0, 1)$.*

A particular case is the following well known result (see e.g. [1, chapter 13]):

Corollary 1.3. *The Riemann zeta function $\zeta(s)$ has no zeros in $(0, 1)$.*

Proof. In fact $\zeta(s) = \zeta(s, 1)$. \square

Remark 1.4. It is well known that the Dirichlet L -function, defined for $\Re(s) > 1$ as $L(s, \chi) \doteq \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$, where χ is a character (mod q), can be represented through a sum of Hurwitz zeta functions (see e.g [2] or [1]) in the following way:

$$L(s, \chi) = q^{-s} \sum_{a=1}^q \chi(a) \zeta(s, a/q).$$

The Extended Riemann Hypothesis conjectures that the Dirichlet L -function $L(s, \chi)$, for a primitive character χ , has no zeros with real part different from $1/2$ in the critical strip; and its strong version says that $L(1/2, \chi)$ is always nonzero too: see also [2, section 10.5.7]).

We leave then open for investigation to see whether our main result could help to find new zero-free regions for this class of functions, for example to show that the Dirichlet L -functions are also nonzero on the real axis between 0 and 1, which would thus establish a weaker version of the Extended Riemann Hypothesis, though stronger in including the point $1/2$.

2 Acknowledgements

Thanks are due in particular to Joachim Rosenthal and Alessandro Cobbe for suggestions and comments.

References

- [1] T. M. Apostol. *Introduction to analytic number theory*. Springer-Verlag, New York, 1976.
- [2] H. Cohen. *Number theory. Vol. II. Analytic and modern tools*, volume 240 of *Graduate Texts in Mathematics*. Springer, New York, 2007.
- [3] H. M. Edwards. *Riemann's zeta function*. Dover Publications Inc., Mineola, NY, 2001. Reprint of the 1974 original.
- [4] R. Spira. Zeros of Hurwitz zeta functions. *Math. Comp.*, 30(136):863–866, 1976.